

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 795 604 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
17.09.1997 Bulletin 1997/38

(51) Int Cl.<sup>6</sup>: **C12N 1/20**  
// A61K35/74

(21) Application number: 97830040.8

(22) Date of filing: 05.02.1997

(84) Designated Contracting States:  
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL**  
**PT SE**

(30) Priority: 11.03.1996 IT MI960468

(71) Applicants:  
• **Cavallero Vesely, Renata Marla Anna**  
I-20123 Milano (IT)  
• **De Simone, Claudio**  
I-00040 Ardea (Roma) (IT)

(72) Inventors:  
• **Cavallero Vesely, Renata Marla Anna**  
20123 Milano (IT)  
• **De Simone, Claudio**  
00040 Ardena (Roma) (IT)

(74) Representative: **Righetti, Giuseppe**  
Bugnion S.p.A.  
Via Carlo Farini, 81  
20159 Milano (IT)

(54) **Strains of bacteria with altered metabolism of bile acids and their use**

(57) Strains of bacteria characterized by exhibiting:  
(a) a 7 $\alpha$ -dehydroxylase activity of less than 50%, and  
(b) a bile acid deconjugation activity of less than 50%,  
and descendants, mutants and derivatives thereof pre-

serving activities (a) and (b); and a pharmaceutical composition comprising one or more of such strains and useful of same for preventing and treating diseases associated with or caused by an altered metabolism of bile acids.

**EP 0 795 604 A2**

## Description

The present invention relates to strains of bacteria and pharmaceutical compositions containing one or more of such strains and use of same for preventing and/or treating diseases associated with or caused by an altered metabolism of bile acids.

Hepatic bile is a pigmented isotonic fluid with an electrolyte composition resembling blood plasma. Major components of bile include water (82 percent), bile acids (12 percent), lecithin and other phospholipids (4 percent), and unesterified cholesterol (0.7 percent). Other constituents include conjugated bilirubin, proteins, electrolytes, mucus and the final products of hepatic transformation of drugs, hormones, etc. The liver production of bile, in basal conditions, is approximately 500-1000 ml/day.

The primary bile acids, cholic acid (CA) and chenodeoxycholic acid (CDCA), are synthesized from cholesterol in the liver, conjugated with glycine or taurine, and excreted into the bile. Secondary bile acids, including deoxycholic acid (DCA) and lithocholic acid (LA), are formed in the colon as bacterial metabolites of the primary bile acids. Other bile acids, called tertiary bile acids (e.g.: ursodeoxycholic acid - UDCA), are formed in the gut following the enzymatic epimerization of -OH groups on sterol rings by the intestinal flora.

In normal bile, the ratio of glycine to taurine conjugates is about 2:1, while in patients with cholestasis, increased concentrations of sulfate and glucuronide conjugates of bile acids are often found.

The intestinal microflora transforms the bile acids into different metabolites. These biotransformations include the hydrolysis of the bond between the bile acid and taurine or glycine, with formation of unconjugated or free bile acids and taurine or glycine. The unconjugated bile acids are therefore made available for the oxidation of the hydroxylic groups in positions C3, C7, and C12 and for the dehydroxylation in positions 7 $\alpha$  and 7 $\beta$ . This latter transformation leads to the formation of the secondary bile acids DCA and LA. The primary bile acids, deconjugated but not transformed, and the secondary biliary acids are reabsorbed from the gut lumen and enter the portal bloodstream, then are taken up by hepatocytes, conjugated with glycine or taurine and resecreted into the bile (enterohepatic circulation).

Normally, the bile acid pool circulates approximately 5 to 10 times daily. Intestinal absorption of the pool is about 95% efficient, so fecal loss of bile acids is in the range of 0.3 to 0.6 g/day. The fecal loss is compensated by an equal daily synthesis.

For this reason the composition of the pool of biliary acids present in the bile is the result of complex interactions occurring between the liver and the microflora enzymes.

Deconjugation activity is a characteristic shared by many bacteria, aerobes and anaerobes, but is particularly common among the obligate anaerobic bacteria, i.e. Bacteroides, Eubacteria, Clostridia, Bifidobacteria, etc. The majority of the bacteria is active against both glycine and taurine conjugates; however, some of them have a certain degree of specificity, depending from the bound amino acid, and the number of hydroxides bound to the steroid nucleus. The free biliary acids obtained following the action of the bacterial hydrolases can undergo the oxidation of the hydroxide groups present at the C3, C7 and C12 positions by the hydroxysteroiddehydrogenase.

The interest in the metabolic disorders of biliary acids comes from the hypothesis that biliary acids and/or metabolites thereof are involved in the pathogenesis of some hepato-biliary and gastroenterologic diseases: biliary dyspepsia, cholelithiasis, acute and chronic hepatopathies, inflammatory diseases of the colon, etc.

Very often in literature the hydrophobicity of the bile acid is correlated with detergency; the secondary bile acids are more hydrophobic than the primary bile acids, the deoxycholic acid (DCA) being actually more detergent than the cholic acid (CA). Therefore an increased concentration of DCA in the bile may involve:

- a) an augmentation of the secretion of cholesterol, with increased saturation index;
- b) a cytotoxic effect on the liver cells.

For this reason a qualitative modification of the bile acids pattern could be a decisive factor, especially in treating the above mentioned pathologies.

Thus, there remains a need for effective bacterial strains or compositions that, by reducing the 7 $\alpha$ -dehydroxylase activity and at the same time deconjugation, can be used for treating and/or preventing diseases associated with metabolic disorders of the biliary acids.

No bacteria strains have been found that are capable of qualitatively modifying the bile acid pattern in a such way.

Accordingly, it is one object of the present invention to provide novel strains of bacteria, in particular gram-positive bacteria, which are useful for treating and/or preventing diseases associated with or caused by a metabolic disorder of biliary acids.

It is another object of the present invention to provide pharmaceutical compositions which contain one or more strains of such bacteria and are useful for treating and/or preventing diseases associated with or caused by a metabolic disorder of biliary acids.

The foregoing and other objects, which will become more apparent during the following detailed description, have

been achieved by the inventors, who have found bacteria strains having a reduced or zero 7 $\alpha$ -dehydroxylase activity and a reduced or zero ability to deconjugate bile acids. This is in contrast with the previous known art. Accordingly, the present invention provides the use of such strains to modify the bile acid metabolism in a useful manner to prevent or treat diseases caused by or associated with metabolic disorders of biliary acids.

Thus, in a first embodiment, the present invention provides novel strains of bacteria which have a 7 $\alpha$ -dehydroxylase activity of less than 50%, preferably less than 25%, and a conjugated bile acid deconjugation activity of less than 50%, preferably less than 25%.

The essential features of the strains according to the present invention are defined in claim 1; specific strains having said features are defined in the dependent claims 2 to 11.

The present invention also provides a pharmaceutical composition for treating and/or preventing diseases associated with or caused by an altered metabolism of biliary acids, said composition comprising at least one bacteria strain according to the present invention. The essential features of the composition according to the present invention are defined in claim 12; specific embodiments of said composition are defined in the dependent claims 13 to 25.

Moreover, the present invention provides the use of at least one bacteria strain according to the present invention to prepare said pharmaceutical composition. The essential features of said use are defined in claim 26; specific embodiments are defined in the dependent claims 27 to 36.

In the context of the present invention, the diseases associated with or caused by a metabolic disorder of the biliary acids include liver diseases and diseases of the digestive apparatus, such as blind loop syndrome, biliary gallstones, cirrhosis, chronic hepatopathies, acute hepatopathies, cystic fibrosis, intrahepatic cholestasis, intestinal inflammatory diseases, colonopathies, malabsorption. The present pharmaceutical compositions may also be used to prevent the onset of biliary gallstones in women during pregnancy or subsequent periods and in subjects undergoing weight-loss programmes or diets.

The 7 $\alpha$ -dehydroxylase activity of the bacteria strain should be less than 50%, preferably less than 25%. The 7 $\alpha$ -dehydroxylase activity values are those measured by the method described in Example 1 below. Specifically, the 10<sup>7</sup> cells of the strain in question are incubated at 37°C for 48 hours, in 15 ml of the specific culture medium with the addition of 2 mg/ml of glycocholic acid (GCA) or 2 mg/ml of taurocholic acid (TCA), and then the amount of 7 $\alpha$ -dehydroxylated product is measured. The percentage value for the 7 $\alpha$ -dehydroxylase activity is calculated by the following formula:

$$7\alpha\text{-dehydroxylase activity} = \frac{\text{mass of } 7\alpha\text{-dehydroxylated product after 48 hours of incubation}}{\text{mass of GCA or TCA at the start of incubation}} \times 100$$

Based on the above, the bacteria strain to be administered should in addition have a conjugated bile acid deconjugation activity of less than 50%, preferably less than 25%. The ability to deconjugate bile acid is determined by using the same incubation procedure described for measuring the 7 $\alpha$ -dehydroxylase activity followed by measuring the amount of deconjugated product formed. The deconjugation activity is calculated using the following formula:

$$\text{Deconjugation activity} = \frac{\text{mass of deconjugated GCA or TCA after 48 hrs of incubation}}{\text{mass of GCA or TCA at the start of incubation}} \times 100$$

The bacteria strains of the present invention may be administered enterically. Preferably, the bacteria strains of the present invention are administered orally.

Although a single bacteria strain may be administered, it is also possible to administer a mixture of two or more bacteria according to the present invention.

Although the exact dosage of bacteria to be administered will vary with the condition and size of the patient, the exact disease being treated, and the identity of the strains being administered, good results have been achieved by administering 10<sup>3</sup> to 10<sup>13</sup> cells of the bacteria/g, preferably 10<sup>8</sup> to 10<sup>12</sup> cells of the bacteria strain/g. To achieve the good effects of the present invention, it is preferred that the strain be administered in an amount and a concentration sufficient to result in the intestines of the patient being populated with an important amount thereof. Thus, it is preferred that the strain be administered in a composition which contains 10<sup>3</sup> to 10<sup>13</sup> cells of the strain/g, preferably 10<sup>8</sup> to 10<sup>12</sup> cells of the strain/g and that the composition be administered in such a regimen so that the patient receives 100 mg to 100 g of the strain/day, preferably 1 g to 20 g of the strain/day, for a period of 1 to 365 days, preferably 3 to 60 days in case of therapy, or according to periodical cycles in case of prophylaxis. The bacteria strain may be administered in any form suitable for enteral administration, such as capsules, tablets, or liquids for oral administration or liquids for enteral administration.

Typically, the administration of the bacteria strain according to the present invention can be prescribed after the diagnosis of metabolic disorders of the biliary acids. However, in the case of the prophylaxis of biliary gallstones, the strain may be administered when the subject is determined to belong to an at-risk population, such as becoming preg-

nant or beginning a weight-loss program or diet. In addition, the present strain of bacteria may be administered after a patient has had their gallbladder removed.

In a preferred embodiment, the coadministration of lactulose is provided when the disease being treated is cirrhosis. Suitably, the lactulose is administered in an amount of 100 mg to 100 g/day, preferably 1 g to 20 g/day.

In another preferred embodiment, the coadministration of bile acid-based preparations, such as ursodeoxycholic acid or tauroursodeoxycholic acid, is provided. Suitably, the ursodeoxycholic or tauroursodeoxycholic acid is administered in an amount of 10 to 3,000 mg/day, preferably 50 to 800 mg/day.

Preferably, the present pharmaceutical compositions contain the strain(s) of bacteria in a concentration of  $10^3$  to  $10^{13}$  cells/g, preferably  $10^8$  to  $10^{12}$  cells/g. The pharmaceutically acceptable carrier may be any which is suitable for enteral administration and is compatible with the strain of bacteria, such as dextrose, calcium carbonate together with different additional substances such as starch, gelatin, vitamins, antioxidants, stains or taste-improving substances.

As an optional component, the compositions of the invention may possibly contain a drug compatible with the bacteria employed and capable of potentiating the activity of the active ingredients present. Anticholinergic drugs, antihistamines, adrenergic, antiulcer, antiacid, antidiarrhoeic and anti-inflammatory drugs, sedatives, antipyretics, chol-  
 15 eretics, anti rheumatic agents, analgesic drugs, diuretics, antiseptic agents, antilipemic, hepatoprotective drugs and drugs active on the gastrointestinal motility (e.g. trimebutine) may be herein mentioned.

When treating cirrhosis, it is preferred that the pharmaceutical composition further comprise lactulose. Suitably, the composition will contain 100 mg to 100 g/day, preferably 1 g to 20 g/day of lactulose. When treating biliary cirrhosis and chronic hepatitis, it is preferred that the pharmaceutical composition comprise bile acid-based preparations, such  
 20 as ursodeoxycholic acid or tauroursodeoxycholic acid. Suitably, the composition will contain 10 to 3,000 mg/day of such bile acid preparations, preferably 50 to 800 mg/day of ursodeoxycholic acid or tauroursodeoxycholic acid.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

## 25 EXAMPLES

### Example 1

30 Strains of the following species have been tested: Streptococcus thermophilus, Streptococcus faecium, Lactobacillus acidophilus, Lactobacillus bulgaricus, Lactobacillus plantarum, Bifidobacterium infantis. Each strain ( $10^7$  CFU) was cultivated in duplicate in specific nutrient broths (15 ml); "CFU" means "colony forming units".

#### List of the employed culture media depending on the different species

35	<u>Bifidobacterium infantis</u> :	MRS + 0.5% glucose (added after sterilization by diluting a 20% sterile solution)
	<u>Streptococcus thermophilus</u> :	M17
	<u>All the remaining strains</u>	MRS

#### Composition of the MRS broth:

40	g/liter	universal peptone	10.0 g
		meat extract	5.0 g
		yeast extract	5.0 g
45		D(+)-glucose	20.0 g
		potassium hydrogen phosphate	2.0 g
		Tween 80	1.0 g
		dibasic ammonium citrate	2.0 g
		sodium acetate	5.0 g
50		magnesium sulfate	0.1 g
		manganous sulfate	0.05 g

Preparation: dissolve 50 g/l in distilled water, sterilized at 121°C for 15 minutes - pH 6.5 + - 0.1 at 25°C

Composition M17 broth (Merck):

g/liter	soybean flour peptone	5.0 g
	meat peptone	2.5 g
	casein peptone	2.5 g
	yeast extract	2.5 g
	meat extract	5.0 g
	O(+)-lactose	5.0 g
	ascorbic acid	0.5 g
	sodium-6-glycerophosphate	19.0 g
	magnesium phosphate	0.25 g

Preparation: dissolve 42.5 g /l in distilled water, sterilized at 121°C for  
15 minutes - pH 7.2 ± 0.1 at 25°

Bifidobacterium infantis was cultivated under anaerobic conditions since it is known that it is an anaerobic bacterium. After 24 hours of incubation at 37°C to each tube was added an amount of bile salt equivalent to 30 mg in order to obtain a final concentration of 2 mg/ml. The bile acids employed are: glycocholic acid (GCA) and taurocholic acid (TCA), obtained from Sigma Chemicals. Each bile acid was added separately to each series of bacterial cultures.

After 48 hours of incubation, isopropanol, 3 ml, was added for 2 minutes. Then it was centrifuged at 400 rpm for 15 minutes and the supernatant was collected (5 ml). The supernatant was kept refrigerated at -30°C until it was analyzed. The percentage of conjugated bile salt present was determined by HPLC (high performance liquid chromatography) utilizing a Gilson apparatus equipped with a detector Diode array mod 1000 and a Spherisorb 5 µm ODS 2 C18 reverse phase column, a mobile phase composed by methanol/buffered phosphate (20 mMol), pH 2.5 in water/ acetonitrile/water (150:60:20:20 by volume), a fluid speed of 0.85 ml/min, at a wavelength of 205 nm; 100 µl of the sample to be tested, dried under nitrogen, were extracted with 100 µl of the mobile phase containing as an internal standard 7 $\alpha$ -OH-12 $\alpha$ -OH-dihydroxy-5 $\beta$ -cholan-3-one (Calbiochem U.S.A.) at a concentration of 2 mg/ml.

The recovery percentage of the bile acid incubated with the bacterial cultures was calculated by the ratio of the area of the bile acid to be detected (GCA or TCA) to the area of the internal standard. When the quantity of the conjugated bile acid found in the bacterial cultures after 48 hours of incubation was less than 50%, thin layer chromatography (TLC) was performed on silica 60 gel plates to detect the presence of CA and DCA, using a mobile phase of cyclohexane/isopropanol/acetic acid (30:10:1 by volume). On every plate, 20 µl of the alcoholic extract of the sample, 20 µl of a solution of CA and DCA, and 20 µl of CA, 20 µl of DCA, were spotted. The plates after development at room temperature, were treated with sulfuric acid and warmed at 145°C until the appearance of the colored spots.

The results of the deconjugation experiments (Table I) show that 5 out of the 16 strains tested with GCA were able to completely deconjugate the bile acid added to the culture, as previously reported in the literature and widely known to all researchers. Surprisingly, ten strains were able to deconjugate GCA but not completely, ranging from 9 to 90 percent (Table I). There was no difference among aerobic and anaerobic bacteria. Two strains, Streptococcus thermophilus YS 52 and Bifidobacterium infantis Bi 6 do not have any deconjugating activity for GCA. The strain YS 52 in addition does not attack the bile acid - taurine bond.

Only one out of the 16 strains tested was able to totally deconjugate the TCA: the Bifidobacterium infantis Bi 6.

The results of the dehydroxylation experiments (Table II) show that only one (Bi 4) out of the 16 strains is able to completely dehydroxylate GCA. Six strains did not dehydroxylate at all: YS 52; SF 2; SF 4; LA 3; LA 10; Bi 6. The other strains were able to dehydroxylate GCA but not completely, ranging from 9% to 90%. As to TCA, seven strains do not dehydroxylate it at all: YS 52; SF 3; LA 3; LA 10; LB 1; LB 7; LB 77. One strain Bi 6 dehydroxylated TCA completely; the other strains dehydroxylated TCA according to varying percentages.



Table I

Percentage of deconjugation of GCA and TCA by bacterial cultures after 48 hours of incubation.			
BACTERIUM	ACCESSION NO.	GCA%	TCA%
<u>Streptococcus thermophilus</u> YS 46	I-1668	9	9
<u>Streptococcus thermophilus</u> YS 48	I-1669	17	11
<u>Streptococcus thermophilus</u> YS 52	I-1670	0	0
<u>Streptococcus faecium</u> SF 2	I-1671	100	3
<u>Streptococcus faecium</u> SF 3		27	0
<u>Streptococcus faecium</u> SF 4		100	12
<u>Lactobacillus acidophilus</u> LA 3		100	80
<u>Lactobacillus acidophilus</u> LA 10	I-1664	100	95
<u>Lactobacillus bulgaricus</u> LB 1		9	0
<u>Lactobacillus bulgaricus</u> LB 3		20	12
<u>Lactobacillus bulgaricus</u> LB 7		14	0
<u>Lactobacillus bulgaricus</u> LB 77	I-1665	20	0
<u>Bifidobacterium infantis</u> Bi 2	I-1666	80	15
<u>Bifidobacterium infantis</u> Bi 3	I-1667	90	10
<u>Bifidobacterium infantis</u> Bi 4		100	26
<u>Bifidobacterium infantis</u> Bi 6		0	100

Table II

Percentage of dehydroxylation of GCA and TCA by bacterial cultures after 48 hours of incubation.			
BACTERIUM	ACCESSION NO.	GCA%	TCA%
<u>Streptococcus thermophilus</u> YS 46	I-1668	9	9
<u>Streptococcus thermophilus</u> YS 48	I-1669	17	11
<u>Streptococcus thermophilus</u> YS 52	I-1670	0	0
<u>Streptococcus faecium</u> SF 2	I-1671	0	3
<u>Streptococcus faecium</u> SF 3		27	0
<u>Streptococcus faecium</u> SF 4		0	12
<u>Lactobacillus acidophilus</u> LA 3		0	0
<u>Lactobacillus acidophilus</u> LA 10	I-1664	0	0
<u>Lactobacillus bulgaricus</u> LB 1		9	0
<u>Lactobacillus bulgaricus</u> LB 3		20	12
<u>Lactobacillus bulgaricus</u> LB 7		14	0
<u>Lactobacillus bulgaricus</u> LB 77	I-1665	20	0
<u>Bifidobacterium infantis</u> Bi 2	I-1666	80	15
<u>Bifidobacterium infantis</u> Bi 3	I-1667	90	10
<u>Bifidobacterium infantis</u> Bi 4		100	26
<u>Bifidobacterium infantis</u> Bi 6		0	100

These strains have been deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the following accession numbers:

<u>Streptococcus thermophilus</u> YS 46	I-1668
<u>Streptococcus thermophilus</u> YS 48	I-1669
<u>Streptococcus thermophilus</u> YS 52	I-1670
<u>Streptococcus faecium</u> SF 3	I-1671
<u>Lactobacillus bulgaricus</u> LB 1	I-1664
<u>Lactobacillus bulgaricus</u> LB 3	I-1665

(continued)

<u>Lactobacillus bulgaricus</u> LB 7	I-1666
<u>Lactobacillus bulgaricus</u> LB 77	I-1667

The following strains are kept and available at the Centro Ricerche Sitia-Yomo S.p.A. - Strada per Merlino, 3 - ZELO BUON PERSICO (Milan) - Italy, distinguished by the below-reported identifiers:

<u>Streptococcus faecium</u> SF 2	SF 2
<u>Streptococcus faecium</u> SF 4	SF 4
<u>Lactobacillus acidophilus</u> LA 3	LA 3
<u>Lactobacillus acidophilus</u> LA 10	LA 10
<u>Bifidobacterium infantis</u> Bi 2	Bi 2
<u>Bifidobacterium infantis</u> Bi 3	Bi 3
<u>Bifidobacterium infantis</u> Bi 4	Bi 4
<u>Bifidobacterium infantis</u> Bi 6	Bi 6

These results demonstrate that the majority of the strains tested by us have a low capability to deconjugate the bile acids and that there are strains that do not deconjugate at all. This observation is surprising in that it has not been known that the lactic acid bacteria deconjugated the biliary salts. Furthermore, it is evident that the enzymes of the strains are selective for the specific bile acid bound on the side chain. In this study, the clearest example is offered by the Bifidobacterium infantis Bi 6. This strain is able not to deconjugate the glycine-conjugated bile acid but is able to totally deconjugate the taurine-conjugated bile acid. Some other strains (LB 1, LB 7, LB 77, SF 3) are unable to deconjugate TCA but are able to deconjugate GCA to a certain extent.

To conclude, strains have been discovered that have a weak or zero capability to deconjugate and dehydroxylate.

#### Example 2

Three healthy volunteers were tested for their content of bile acids following treatment with a lactobacilli preparation containing  $1 \times 10^{11}$  cells of Streptococcus thermophilus YS 52 per gram for a daily total of 6 g for 28 days. Before beginning the treatment and after 12 hours starvation, the subjects were intubated and the gallbladder bile, following stimulation with ceruletide, was collected and frozen at  $-80^{\circ}\text{C}$ . The gallbladder contraction was assessed by echography and the position of the tube, in the second portion of the duodenum was checked by Rx (fluoroscopy).

After a 4 week treatment, the subjects underwent a second intubation and collection of bile. The bile samples were then tested for their content of some bile acids as previously described. The results are shown in Table III.

Table III

Bile Acid	Patient # 1		Patient # 2		Patient # 3	
	before	after	before	after	before	after
Glychenodeoxycholic	32	15	22	15	28	12
Glycodeoxycholic	6	5	9	2	4	3
Glycoursodeoxycholic	1	5	1	7	1	4
Taurocholic	9	26	15	25	12	21
Taurodeoxycholic	1	3	5	8	3	9
NOTE: (the bile acids are listed following the hydrophilic capacity order, that is in inverse relation to detergency)						
Taurocholic						
Taurodeoxycholic						
Glycoursodeoxycholic						
Glycodeoxycholic						
Glychenodeoxycholic						



This experiment is a confirmation of what shown in Example No. 1, that is a lower deconjugation in one of the primary bile acids if bacteria being the object of the present invention are administered. The achieved result is a longer maintenance of the primary bile acids in the enterohepatic circulation.

The properties of the bile acids are reported in the note to Table III. Thus, according to these results the administration of selected strains of bacteria can reduce the detergency property and therefore the cytolytic activity of bile acids.

### Example 3

Fourteen patients with chronic hepatitis were treated with a bacterial preparation containing Streptococcus thermophilus YS 46 and YS 48 (two strains), Lactobacillus bulgaricus LB1, LB7 and LB 77 (three strains). Each strain had been brought to a concentration of  $150 \times 10^9$  cells per gram before being mixed with the others, to prepare a mixture containing the same parts by weight of each strain. 6 grams per day of said mixture were administered for 28 days. Liver enzymes were measured before and after the treatment, and the results are shown in Table IV.

Table IV

Influence of the Treatment with the Bacterial Mixture on Liver Enzymes Aspartate Transaminase (AST; SGOT) and alanine transaminase (ALT; SGPT)				
Patient	AST (SGOT)		ALT (SGPT)	
	before	after	before	after
# 1	92	59	102	46
# 2	89	67	96	42
# 3	174	86	97	39
# 4	121	91	102	66
# 5	116	81	111	55
# 6	156	87	94	76
# 7	163	66	69	37
# 8	78	64	122	57
# 9	109	39	87	86
# 10	166	70	102	48
# 11	56	24	118	62
# 12	131	83	96	79
# 13	137	86	94	74
# 14	84	87	144	114
Mean	119	71	102	63
Standard deviation	36	19	17	21
Significance Student t test for paired data	p<0,001		p<0,001	

### Claims

1. A bacteria strain characterized by exhibiting:
  - (a) a 7 $\alpha$ -dehydroxylase activity of less than 50%, and
  - (b) a bile acid deconjugation activity of less than 50%, and descendants, mutants and derivatives thereof preserving activities (a) and (b).
2. The strain of claim 1, which is a gram-positive bacteria strain.
3. The strain of claim 1 and/or 2, belonging to a species selected from Streptococcus thermophilus, Streptococcus faecium and Lactobacillus bulgaricus.
4. The strain of claim 3, wherein the bacteria strain is Streptococcus thermophilus YS 52 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1670.

5. The strain of claim 3, wherein the bacteria strain is Streptococcus thermophilus YS 46 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1668.
6. The strain of claim 3, wherein the bacteria strain is Streptococcus thermophilus YS 48 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1669.
7. The strain of claim 3, wherein the bacteria strain is Streptococcus faecium SF 3 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1671.
8. The strain of claim 3, wherein the bacteria strain is Lactobacillus bulgaricus LB 1 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1664.
9. The strain of claim 3, wherein the bacteria strain is Lactobacillus bulgaricus LB 3 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1665.
10. The strain of claim 3, wherein the bacteria strain is Lactobacillus bulgaricus LB 7 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1666.
11. The strain of claim 3, wherein the bacteria strain is Lactobacillus bulgaricus LB 77 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1667.
12. A pharmaceutical composition for preventing and/or treating diseases associated with or caused by an altered metabolism of biliary acids, characterized in that it comprises an effective amount capable of producing a normalizing effect on such an altered metabolism in a patient suffering therefrom, of
  - (1) at least one bacteria strain provided with:
    - (a) a 7 $\alpha$ -dehydroxylase activity of less than 50%, and
    - (b) a bile acid deconjugation activity of less than 50%, and descendants, mutants and derivatives thereof preserving activities (a) and (b), and
  - (2) a pharmaceutically acceptable carrier.
13. The pharmaceutical composition of claim 12, wherein said at least one bacteria strain is a gram-positive bacteria strain.
14. The composition of claim 12 and/or 13, wherein said at least one bacteria strain belongs to a species selected from the group consisting of Streptococcus thermophilus, Streptococcus faecium and Lactobacillus bulgaricus.
15. The composition of claim 14, wherein the bacteria strain is Streptococcus thermophilus YS 52 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1670.
16. The composition of claim 14, wherein the bacteria strain is Streptococcus thermophilus YS 46 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1668.
17. The composition of claim 14, wherein the bacteria strain is Streptococcus thermophilus YS 48 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1669.
18. The composition of claim 14, wherein the bacteria strain is Streptococcus faecium SF 3 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1671.
19. The composition of claim 14, wherein the bacteria strain is Lactobacillus bulgaricus LB 1 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1664.
20. The composition of claim 14, wherein the bacteria strain is Lactobacillus bulgaricus LB 3 deposited with the CNCM

- Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1665.

21. The composition of claim 14, wherein the bacteria strain is Lactobacillus bulgaricus LB 7 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1666.

22. The composition of claim 14, wherein the bacteria strain is Lactobacillus bulgaricus LB 77 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1667.

23. The composition of claim 12, comprising  $10^3$  to  $10^{13}$  cells of the bacteria strain per gram of composition.

24. The composition of claim 12, further comprising lactulose.

25. The composition of claim 12, further comprising bile acid-based preparations selected from ursodeoxycholic acid and tauroursodeoxycholic acid.

26. Use of at least one bacteria strain characterized by exhibiting:

(a) a 7 $\alpha$ -dehydroxylase activity of less than 50%, and

(b) a bile acid deconjugation activity of less than 50%, and descendants, mutants and derivatives thereof preserving activities (a) and (b), to prepare a pharmaceutical composition for preventing and treating diseases caused by or associated with an altered metabolism of the bile acids.

27. The use of claim 26, wherein the at least one bacteria strain is a gram-positive bacteria strain.

28. The use of claim 26 and/or 27, wherein the bacteria strain belongs to a species selected from the group consisting of Streptococcus thermophilus, Streptococcus faecium and Lactobacillus bulgaricus.

29. The use of claim 28, wherein the bacteria strain is Streptococcus thermophilus YS 52 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1670.

30. The use of claim 28, wherein the bacteria strain is Streptococcus thermophilus YS 46 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1668.

31. The use of claim 28, wherein the bacteria strain is Streptococcus thermophilus YS 48 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1669.

32. The use of claim 28, wherein the bacteria strain is Streptococcus faecium SF 3 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1671.

33. The use of claim 28, wherein the bacteria strain is Lactobacillus bulgaricus LB 1 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1664.

34. The use of claim 28, wherein the bacteria strain is Lactobacillus bulgaricus LB 3 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1665.

35. The use of claim 28, wherein the bacteria strain is Lactobacillus bulgaricus LB 7 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1666.

36. The use of claim 28, wherein the bacteria strain is Lactobacillus bulgaricus LB 77 deposited with the CNCM - Collection Nationale de Cultures de Microorganismes - Institut Pasteur, under the accession number I-1667.